

On classification of projective homogeneous varieties up to motivic isomorphism

N. Semenov, K. Zainoulline

February 2, 2008

Abstract

We give a complete classification of anisotropic projective homogeneous varieties of dimension less than 6 up to motivic isomorphism. We give several criteria for anisotropic flag varieties of type A_n to have isomorphic motives.

1 Introduction

The present paper can be viewed as an application of the methods and results obtained by the authors in [CPSZ].

Let k be a field of characteristic not 2 and k_s denotes its separable closure. For a variety X over k we denote by X_s the base change $X \times_k k_s$. By $\mathcal{M}(X)$ we denote the Chow motive of X . Recall (see [MPW, § 1]) that X is a twisted flag variety of inner type over k if $X = {}_\xi(G/P)$ is a twisted form of the projective homogeneous variety G/P , where G is an adjoint simple split algebraic group over k , P its parabolic subgroup and the twisting is given by a 1-cocycle $\xi \in Z^1(k, G(k_s))$.

The present paper is devoted to the following

Problem. *Describe all pairs (X, Y) of non-isomorphic twisted flag varieties X and Y of inner type over k which have isomorphic Chow motives.*

This problem can be subdivided into two subproblems:

- (i) Describe all such pairs (X, Y) with $X_s \simeq Y_s$;

(ii) Describe all such pairs (X, Y) with $X_s \not\simeq Y_s$.

Let us briefly remind what is known so far. The complete solution of the problem (i) is known for quadrics and Severi-Brauer varieties due to Izhboldin, Karpenko, Merkurjev, Rost, Vishik and others (see [Izh98], [Ka96], [Ka00], [Ro98], [Vi03]). Concerning (ii), the example (of dimension 5) was provided by Bonnet in [Bo03]. It deals with twisted flag varieties of type G_2 . For exceptional varieties of type F_4 a similar example was provided in [NSZ].

In the present paper we provide a complete solution of the mentioned above problem for projective homogeneous varieties of dimension less than 6. Namely, we prove the following (using the notation of 2.1)

1.1 Theorem. *Let X and Y be non-isomorphic twisted flag varieties of dimension ≤ 5 of inner type over k which have isomorphic Chow motives.*

(i) *If $X_s \simeq Y_s$, then either*

$X = \text{SB}(A)$ and $Y = \text{SB}(A^{\text{op}})$ are Severi-Brauer varieties corresponding to a central simple algebra A and its opposite A^{op} , where $\deg(A) = 3, 4, 5, 6$ and $\exp(A) > 2$

or

$X = \text{SB}_{1,2}(A)$ and $Y = \text{SB}_{1,2}(A^{\text{op}})$ are twisted forms of the flag varieties corresponding to central simple algebras A such that $\deg(A) = \exp(A) = 4$.

(ii) *If $X_s \not\simeq Y_s$, then either*

$X = \text{SB}_{1,3}(A)$ and $Y = \text{SB}_{1,2}(A')$ are twisted forms of the flag varieties corresponding to central simple algebras A and A' such that $\deg(A) = \deg(A') = 4$ and $A \simeq A'$ or A'^{op} ,

or

$X = {}_{\xi}(G_2/P_1)$ and $Y = {}_{\xi}(G_2/P_2)$ are twisted forms of the variety G/P_i , $i = 1, 2$, where G is a split exceptional group of type G_2 and P_i is one of its maximal parabolic subgroups,

or

$X = \mathbb{P}^n$ and $Y = Q^n$ is the projective space and the split quadric respectively, where $n = 3, 5$.

or

$X = \mathbb{P}^5$ and $Y = G_2/P_2$ is the projective space and the split Fano variety of type G_2 .

1.2 Remark. Observe that the case $X = {}_{\xi}(G_2/P_1)$ and $Y = {}_{\xi}(G_2/P_2)$ of the theorem is the example of Bonnet mentioned above and, hence, is the minimal one in the sense of dimension.

1.3 Remark. The case $X = \text{SB}_{1,3}(A)$ and $Y = \text{SB}_{2,3}(A')$ with $A \simeq A', A'^{\text{op}}$ provides another minimal example of two non-isomorphic varieties that have isomorphic Chow motives.

Apart from Theorem 1.1, we prove the following

1.4 Theorem. *Let $X = \text{SB}_{n_1, \dots, n_r}(A)$ and $Y = \text{SB}_{m_1, \dots, m_r}(A')$ be twisted flag varieties of inner type A_n , $n \geq 2$, over k , where the central simple algebras A and A' have exponents 1, 2, 3, 4, or 6. Assume that*

- (i) $\mathcal{M}(X_s) \simeq \mathcal{M}(Y_s)$;
- (ii) $n_1 = 1$ or $n_r = n$
- (iii) $m_1 = 1$ or $m_r = n$.

Then $\mathcal{M}(X) \simeq \mathcal{M}(Y) \Leftrightarrow A \simeq A'$ or A'^{op} .

As an immediate consequence we obtain

1.5 Corollary. *Let $X = \text{SB}_{1,n}(A)$ and $Y = \text{SB}_{n-1,n}(A')$, where A and A' are central simple algebras of degree $n+1$, $n \geq 3$, and exponent 1, 2, 3, 4 or 6. Then*

$$\mathcal{M}(X) \simeq \mathcal{M}(Y) \Leftrightarrow A \simeq A' \text{ or } A'^{\text{op}}.$$

1.6 Remark. The varieties X and Y of 1.5 provide examples of twisted flag varieties that satisfy **(b)**. In fact, $X_s \not\simeq Y_s$, since they have different automorphism groups by [De77].

The paper is organized as follows. In section 2 we consider the case of a split group and state several facts which will be extensively used in the proofs. Section 3 is devoted to the case by case proof of Theorem 1.1. In the section 4 we prove Theorem 1.4 and provide several results that we need for the proof of 1.1.

2 Preliminaries

In the paper we use the following notation.

2.1. Let G be a split simple algebraic group defined over a field k . We fix a maximal split torus T of G and a Borel subgroup B of G containing T and defined over k . Denote by Φ the root system of G , by $\Pi = \{\alpha_1, \dots, \alpha_{\text{rk } G}\}$ the set of simple roots of Φ corresponding to B , by W the Weyl group, and by $S = \{s_1, \dots, s_{\text{rk } G}\}$ the corresponding set of fundamental reflections. Let P_Θ be the standard parabolic subgroup corresponding to a subset $\Theta \subset \Pi$, i.e., $P_\Theta = BW_\Theta B$, where $W_\Theta = \langle s_\theta, \theta \in \Theta \rangle$. Denote by P_i the maximal parabolic subgroup $P_{\Pi \setminus \{\alpha_i\}}$. By Φ/P_Θ we denote the flag variety G/P_Θ . The root enumeration follows Bourbaki.

The notation $\text{SB}_{n_1, \dots, n_r}(A)$, $1 \leq n_1 < \dots < n_r \leq n$, is used for the twisted form of the variety A_n/P_Θ , where $\Theta = \Pi \setminus \{\alpha_{n_1}, \dots, \alpha_{n_r}\}$ and A is a central simple algebra of degree $n+1$ corresponding to the twisting. Observe that $\text{SB}_{n_1, \dots, n_r}(A) = X(A; n_1, \dots, n_r)$ in the notation of [MPW, Appendix] and $\text{SB}(A) = \text{SB}_1(A)$ is the usual Severi-Brauer variety defined by A . By $\text{ind}(A)$ we denote the index of A and by $\text{exp}(A)$ its exponent. A split projective quadric of dimension n is denoted by Q^n .

2.2. According to [Ko91] the Chow motive of the flag variety $X = G/P_\Theta$, when G is a split group, is isomorphic to

$$\mathcal{M}(X) \simeq \bigoplus_{i=0}^{\dim X} \mathbb{Z}(i)^{\oplus a_i(X)},$$

where $\mathbb{Z}(i)$ are the twists of the Lefschetz motive and the positive integers (ranks) $a_i(X)$ are the coefficients of the generating polynomial $p_X(z) = \sum_{i=0}^{\dim X} a_i(X) z^i$. The latter is defined by the following explicit formula:

$$p_X(z) = \left(\prod_{i=1}^{\text{rk } G} \frac{z^{d_i(W)} - 1}{z - 1} \right) / \left(\prod_{j=1}^m \prod_i \frac{z^{d_i(W_j)} - 1}{z - 1} \right).$$

Here $W_1 \times \dots \times W_m$ is the decomposition of W_Θ into the product of Weyl groups corresponding to the irreducible root systems and $d_i(W_j)$ are the degrees of the respective fundamental polynomial invariants (see [Ca72, 9.4 A]).

The following observation follows from the above isomorphism.

2.3. *The motives of flag varieties X and Y of dimension n over a separably closed field are isomorphic iff the corresponding sequences of ranks $(a_0(X), \dots, a_n(X))$ and $(a_0(Y), \dots, a_n(Y))$ are equal.*

We shall need the following two facts:

2.4. (See [Ka00, Criterion 7.1]) Let A, A' be central simple algebras over k and $\text{SB}(A), \text{SB}(A')$ be the respective Severi-Brauer varieties. Then

$$\mathcal{M}(\text{SB}(A)) \simeq \mathcal{M}(\text{SB}(A')) \Leftrightarrow A \simeq A', A'^{\text{op}}.$$

2.5. (see [Izh98, Cor. 2.9 and Prop. 3.1]) Let q, q' be regular quadratic forms of rank n and $X_q, X_{q'}$ be the respective projective quadrics. If n is odd or $n < 7$, then

$$\mathcal{M}(X_q) \simeq \mathcal{M}(X_{q'}) \Leftrightarrow X_q \simeq X_{q'}.$$

Finally, we shall need the following observation:

2.6. (See [Ka00, Proof of Lemma 2.3]) Let X and Y be smooth projective varieties over k with isomorphic Chow motives. Then there is an isomorphism of abelian groups

$$\text{Coker}(\text{CH}_0(X) \xrightarrow{\text{res}} \text{CH}_0(X_s)) \simeq \text{Coker}(\text{CH}_0(Y) \xrightarrow{\text{res}} \text{CH}_0(Y_s)).$$

3 Small dimensions

In this section we classify all pairs (X, Y) of non-isomorphic twisted flag varieties of inner type over k of dimension ≤ 5 which have isomorphic Chow motives and hence prove Theorem 1.1.

Dimension 1. Twisted flag varieties of dimension 1 are the twisted forms of the projective line $\mathbb{P}^1$. The twisted forms of $\mathbb{P}^1$ are Severi-Brauer varieties $\text{SB}(H)$, where H is a quaternion algebra. By 2.4

$$\mathcal{M}(\text{SB}(H)) \simeq \mathcal{M}(\text{SB}(H')) \Leftrightarrow H \simeq H', H'^{\text{op}}$$

Since $H \simeq H'^{\text{op}}$, we conclude that the motives are isomorphic iff the varieties are isomorphic.

Dimension 2. All twisted flags of dimension 2 are the twisted forms of the projective space $\mathbb{P}^2$ or the split quadric surface $Q^2 \simeq \mathbb{P}^1 \times \mathbb{P}^1$. Observe that Q^2 is a projective homogeneous variety for a group of type D_2 which is not simple, but semisimple. Nevertheless, we shall consider this case too.

The motives of $\mathbb{P}^2$ and Q^2 are not isomorphic, since the respective sequences of ranks $(1, 1, 1)$ and $(1, 2, 1)$ are different.

The twisted forms of Q^2 of inner type over k are 2-dimensional quadrics (see [Inv, Cor. (15.12)]). By 2.5 the motives of two quadrics of dimension 2 are isomorphic iff the quadrics are isomorphic.

The twisted forms of $\mathbb{P}^2$ are Severi-Brauer varieties $SB(A)$, where A is a central simple algebra of degree 3. Again by 2.4 we have

$$\mathcal{M}(SB(A)) \simeq \mathcal{M}(SB(A')) \Leftrightarrow A \simeq A', A'^{\text{op}}.$$

Since the varieties $SB(A)$ and $SB(A^{\text{op}})$ are isomorphic iff A is split, we conclude that all pairs of non-isomorphic varieties which have isomorphic motives are of the kind $(SB(A), SB(A^{\text{op}}))$, where A is a division algebra of degree 3.

Dimension 3. Computing generating functions (see 2.2) we conclude that there are only three projective homogeneous varieties of dimension 3 over k_s . Namely, the projective space $\mathbb{P}^3$, the quadric Q^3 and the variety of complete flags A_2/B (B denotes a Borel subgroup). The respective sequences of ranks look as follows:

$$\begin{array}{ll} \mathbb{P}^3 \simeq A_3/P_1 & : (1, 1, 1, 1) \\ Q^3 \simeq B_2/P_1 & : (1, 1, 1, 1) \\ A_2/B & : (1, 2, 2, 1) \end{array}$$

In particular, we see that the motives of $\mathbb{P}^3$ and Q^3 are isomorphic but the motives of Q^3 and A_2/B are not.

By 2.4 all non-isomorphic twisted forms of $\mathbb{P}^3$ which have isomorphic motives form pairs $(SB(A), SB(A^{\text{op}}))$, where A is a division algebra of degree 4 and exponent 4. Observe that all non-isomorphic twisted forms of Q^3 are quadrics as well and by 2.5 the motive of a quadric determines this quadric uniquely. Therefore it remains to describe all possible motivic isomorphisms between the twisted forms ${}_{\xi}\mathbb{P}^3$ and ${}_{\zeta}Q^3$ and the twisted forms ${}_{\xi}(A_2/B)$ and ${}_{\zeta}(A_2/B)$ of the variety of complete flags A_2/B .

According to Corollary 4.4 there are no non-isomorphic twisted forms of A_2/B which have isomorphic Chow motives. And the next lemma shows that there are no such (non-trivial) twisted forms of $\mathbb{P}^3$ and Q^3 .

3.1 Lemma. *Let ξ, ζ be 1-cocycles. Then $\mathcal{M}(\xi\mathbb{P}^3) \simeq \mathcal{M}(\zeta Q^3)$ iff ξ and ζ are trivial.*

Proof. This is a particular case of a more general result (see Lemma 4.2) proven using Index Reduction Formula. Here we give an elementary proof. It uses only well-known facts about quadrics and Severi-Brauer varieties.

Observe that any twisted form of $\mathbb{P}^3$ is a Severi-Brauer variety $\text{SB}(A)$ for some central simple algebra A of degree 4 and any twisted form of Q^3 is a non-singular quadric of dimension 3.

As in 2.6 for a variety X consider the abelian group $\text{Coker}(\text{CH}_0(X) \rightarrow \text{CH}_0(X_s))$. If $X = \text{SB}(A)$ is a Severi-Brauer variety of a central simple algebra A , then this cokernel is equal to $\mathbb{Z}/\text{ind}(A)\mathbb{Z}$ (see [Ka00]), where $\text{ind}(A)$ is the index of A . In particular, this cokernel is trivial iff A is split. If X is a quadric then this cokernel is trivial iff X is isotropic (see [Sw89]). In the case X is an anisotropic quadric this cokernel is isomorphic to $\mathbb{Z}/2\mathbb{Z}$.

In our case we have two varieties $X = \text{SB}(A)$ and $Y = {}_\zeta Q^3$ which have isomorphic motives. Hence, by 2.6 the respective cokernels must be isomorphic.

Hence, if the quadric Y is isotropic, then the algebra A is split. The latter implies that the motive $\mathcal{M}(\text{SB}(A))$ splits into the direct sum of Lefschetz motives and so is $\mathcal{M}(Y)$, i.e., Y is split as well by 2.5.

Assume q is anisotropic, then there exists a quadratic field extension l/k such that the Witt index of $Y_l = Y \times_k l$ is one (see [Vi03, §7.2]). Since the motives of X and Y are still isomorphic over l , we conclude that A is split over l . Then Y_l is split as well. This leads to a contradiction. $\square$

3.2 Remark. Observe that the pair of twisted forms $(\xi(\text{B}_2/P_1), \xi(\text{B}_2/P_2))$ can be viewed as a low-dimensional analog of the pair $(\xi(\text{G}_2/P_1), \xi(\text{G}_2/P_2))$ considered by Bonnet. The lemma says that contrary to G_2 -case the motives of $\xi(\text{B}_2/P_1)$ and $\xi(\text{B}_2/P_2)$ are not isomorphic (if ξ is non-trivial).

Dimension 4. There are three non-isomorphic projective homogeneous varieties of dimension 4 over k_s . Namely, the projective space $\mathbb{P}^4$, the 4-dimensional quadric $Q^4 \simeq \text{Gr}(2, 4)$ and the variety of complete flags B_2/B . The respective sequences of ranks in these cases are all different and look as follows:

$$\begin{array}{ll} \mathbb{P}^4 \simeq \text{A}_4/P_1 & : (1, 1, 1, 1, 1) \\ Q^4 \simeq \text{A}_3/P_2 & : (1, 1, 2, 1, 1) \\ \text{B}_2/B & : (1, 2, 2, 2, 1) \end{array}$$

Hence, the motives of $\mathbb{P}^4$, Q^4 and B_2/B are non-isomorphic to each other.

By 2.4 all non-isomorphic twisted forms of $\mathbb{P}^4$ which have isomorphic motives form pairs $(SB(A), SB(A^{\text{op}}))$, where A is a division algebra of degree 5. By Corollary 4.5 there are no non-isomorphic twisted forms of B_2/B which have isomorphic Chow motives. Therefore the only case left is the case of inner twisted forms of Q^4 .

The inner forms of Q^4 are the generalized Severi-Brauer varieties $SB_2(A)$, where A is a central simple algebra of degree 4. The next lemma shows that there are no non-isomorphic forms of $SB_2(A)$ which have isomorphic motives.

3.3 Lemma. *Let A, A' be central simple algebras of degree 4. Then*

$$\mathcal{M}(SB_2(A)) \simeq \mathcal{M}(SB_2(A')) \Leftrightarrow SB_2(A) \simeq SB_2(A')$$

Proof. Let $\mathcal{M}(SB_2(A)) \simeq \mathcal{M}(SB_2(A'))$. It suffices to prove that for all field extensions l/k one has $\text{ind}(A_l) = \text{ind}(A'_l)$. Indeed, by [Ka00, Lemma 7.13] $\langle A \rangle = \langle A' \rangle$ in $\text{Br}(k)$, hence, $A \simeq A'$ or A^{op} . But $SB_2(A) \simeq SB_2(A^{\text{op}})$ for any central simple algebra A of degree 4.

Assume that there exists a field extension l/k such that $\text{ind}(A_l) \neq \text{ind}(A'_l)$. Depending on the indices of A and A' we distinguish the following cases:

Case 1. $\text{ind}(A) = 4$ and $\text{ind}(A') = 1$ or 2 .

In this case $SB_2(A')$ has a rational point. By [Inv, Case $A_3 = D_3$], the variety $SB_2(A')$ is isotropic, hence, the group

$$\text{Coker}(\text{CH}_0(SB_2(A')) \rightarrow \text{CH}_0(SB_2(A'_{k_s})))$$

is trivial. By 2.6 the cokernel

$$\text{Coker}(\text{CH}_0(SB_2(A)) \rightarrow \text{CH}_0(SB_2(A_{k_s})))$$

must be trivial as well. If $\exp(A) = 2$, then A is a biquaternion algebra and by [Inv, Cor. (15.33)] $SB_2(A)$ is an anisotropic quadric. Then the cokernel above must be isomorphic to $\mathbb{Z}/2\mathbb{Z}$, a contradiction. If $\exp(A) = 4$, then by [Inv, Cor. (15.33)] $A \simeq C^\pm(B, \sigma, f)$, where $(B, \sigma, f) \in {}^1D_3$ and B is a central simple algebra of degree 6 and index 2. By Merkurjev's theorem (see [Me95]) the cokernel above must be again isomorphic to $\mathbb{Z}/2\mathbb{Z}$, a contradiction.

Case 2. $\text{ind}(A) = 2$ and $\text{ind}(A') = 1$.

In this case A' is split, hence, the corresponding variety is a split quadric. From the other hand, $\text{SB}_2(A) \simeq X_q$, where q is some 6-dimensional quadratic form and X_q is the corresponding projective quadric. Using 2.5, we conclude that $\text{SB}_2(A) \simeq \text{SB}_2(A')$, a contradiction. $\square$

Dimension 5. There are five non-isomorphic projective homogeneous varieties over k_s of dimension 5. Namely, the projective space $\mathbb{P}^5$, the quadric Q^5 , the exceptional Fano variety G_2/P_2 , the flag varieties $A_3/P_{\{\alpha_1\}}$ and $A_3/P_{\{\alpha_2\}}$. The respective sequences of ranks look as follows:

$$\begin{array}{ll} \mathbb{P}^5 \simeq A_5/P_1 & : (1, 1, 1, 1, 1, 1) \\ Q^5 \simeq B_3/P_1 & : (1, 1, 1, 1, 1, 1) \\ G_2/P_2 & : (1, 1, 1, 1, 1, 1) \\ A_3/P_{\{\alpha_1\}} \simeq A_3/P_{\{\alpha_3\}} & : (1, 2, 3, 3, 2, 1) \\ A_3/P_{\{\alpha_2\}} & : (1, 2, 3, 3, 2, 1) \end{array}$$

Therefore, the motives of $\mathbb{P}^5$, Q^5 and G_2/P_2 are isomorphic and the motives of $A_3/P_{\{\alpha_1\}}$ and $A_3/P_{\{\alpha_2\}}$ are isomorphic.

As was mentioned before, the twisted forms of $\mathbb{P}^5$ and Q^5 were completely classified up to motivic isomorphisms by Karpenko and Izhiboldin (see 2.4 and 2.5). Moreover, by Lemma 4.2 there is only one pair $(\xi\mathbb{P}^5, \zeta Q^5)$ of twisted forms that have isomorphic motives.

By the result of Bonnet [Bo03] the motive of the twisted form $\xi(G_2/P_2)$ is isomorphic to the motive of $\xi(G_2/P_1)$ which is a 5-dimensional quadric.

By Corollary 1.5 the motives of the twisted forms of $A_3/P_{\{\alpha_1\}}$ and $A_3/P_{\{\alpha_2\}}$ are isomorphic iff the respective central simple algebras of degree 4 are isomorphic or opposite. This provides the last example (see 1.1) of a pair of non-isomorphic varieties of dimension 5 that have isomorphic motives.

4 Arbitrary dimensions

In the present section we prove several classification results. We start with the following

4.1 Lemma. *Let X and Y be twisted flag varieties of inner type over k which have isomorphic Chow motives. Assume X is not of type E_8 and splits over its function field $k(X)$, i.e., the group corresponding to X splits over $k(X)$. Then X splits over the function field of Y .*

Proof. Since the motives are isomorphic, there is an isomorphism of cokernels (see 2.6) and, hence, an isomorphism of cokernels over $k(Y)$

$$\text{Coker}(\text{CH}_0(X_{k(Y)}) \rightarrow \text{CH}_0(X_{k(Y)_s})) \simeq \text{Coker}(\text{CH}_0(Y_{k(Y)}) \rightarrow \text{CH}_0(Y_{k(Y)_s}))$$

Since $Y_{k(Y)}$ is isotropic, the right cokernel is trivial and so is the left one. The fact that the map $\text{res} : \text{CH}_0(X_{k(Y)}) \rightarrow \text{CH}_0(X_{k(Y)_s})$ is surjective and the group $\text{CH}_0(X_{k(Y)_s})$ is a free abelian group of rank one generated by the class of a rational point $[pt]$ implies that the preimage $\text{res}^{-1}([pt])$ is a 0-cycle of degree 1 in $\text{CH}_0(X_{k(Y)})$. Then, the variety $X_{k(Y)}$ is isotropic (see [To04, Q. 0.2]).

By [KR94, 3.16.(iii)] the function field $k(X)$ is a generic splitting field for the respective parabolic subgroup P . Since $X_{k(Y)}$ is isotropic, the field $k(Y)$ is a k -specialization of $k(X)$ (see [KR94, Def. 1.2]). Since X splits over $k(X)$, $k(X)$ is a splitting field for the respective group G . Then, by [KR94, 3.9.(iii)], $k(Y)$ is a splitting field of G as well, i.e., $X_{k(Y)}$ splits. $\square$

4.2 Proposition. *Let γ, δ be 1-cocycles and $X = {}_\gamma\mathbb{P}^n$, $Y = {}_\delta\mathbb{Q}^n$ be the respective twisted forms for $n > 1$ odd. Then*

$$\mathcal{M}(X) \simeq \mathcal{M}(Y) \Leftrightarrow \gamma \text{ and } \delta \text{ are trivial.}$$

Proof. Observe that X is a Severi-Brauer variety corresponding to a central simple algebra A and Y is a n -dimensional quadric.

Assume that $\mathcal{M}(X) \simeq \mathcal{M}(Y)$ and γ is not trivial. By Lemma 4.1 applied to X and Y , the algebra $A_{k(Y)}$ splits, i.e., $\text{ind}(A_{k(Y)}) = 1$. From the other hand by Index Reduction Formula (see [MPW]) we obtain

$$\text{ind}(A_{k(Y)}) = \min\{\text{ind}(A), 2^{(n-1)/2} \text{ind}(A \otimes_k C_0(q))\} > 1,$$

where $C_0(q)$ is the even part of the Clifford algebra of the quadric corresponding to Y . This leads to a contradiction. $\square$

Note that the same proof works for twisted forms of types B_n and C_n . Namely,

4.3 Proposition. *Let γ, δ be 1-cocycles and $X = {}_\gamma(C_n/P_l)$, $Y = {}_\delta(B_n/P_l)$ be the respective twisted forms for an odd $1 \leq l < n$. Then*

$$\mathcal{M}(X) \simeq \mathcal{M}(Y) \Leftrightarrow \gamma \text{ and } \delta \text{ are trivial.}$$

Proof. For any simple algebraic group G as above consider a twisted flag variety $W = {}_{\xi}(G/P_{\Theta})$ over k . On the Tits diagram (see [Ti66]) of G over $k(W)$ all vertices corresponding to simple roots from $\Pi \setminus \Theta$ are circled.

In our case since l is odd, this implies that $X_{k(X)}$ is split (see [Ti66] for a complete list of Tits diagrams). The rest of the proof repeats the proof of 4.2. $\square$

The rest of this section is devoted to the twisted forms of flag varieties. In particular, we obtain the description of motivic isomorphisms for twisted forms of the flag varieties A_2/B , B_2/B and $A_3/P_{\{\alpha_i\}}$, $i = 1, 2, 3$. We start with the proof of Theorem 1.4.

Proof of Theorem 1.4. Assume $\mathcal{M}(X) \simeq \mathcal{M}(Y)$. Since X and Y are twisted forms of flags containing the subspace of dimension 1 (we may assume $n_1 = m_1 = 1$), the motives of X and Y can be decomposed into a direct sum of twisted motives of Severi-Brauer varieties (see [CPSZ, Thm. 2.1]). Namely,

$$\mathcal{M}(X) \simeq \bigoplus_i \mathcal{M}(\text{SB}(A))(i), \quad \mathcal{M}(Y) \simeq \bigoplus_j \mathcal{M}(\text{SB}(A'))(j). \quad (*)$$

This together with 2.6 implies the isomorphism of abelian groups

$$\text{Coker}(\text{CH}_0(\text{SB}(A)) \rightarrow \text{CH}_0(\mathbb{P}^n)) \simeq \text{Coker}(\text{CH}_0(\text{SB}(A')) \rightarrow \text{CH}_0(\mathbb{P}^n))$$

and, hence, the isomorphism $\mathbb{Z}/\text{ind}(A)\mathbb{Z} \simeq \mathbb{Z}/\text{ind}(A')\mathbb{Z}$, i.e., $\text{ind}(A) = \text{ind}(A')$. Since the motivic isomorphism is preserved under the base extensions, we obtain that $\text{ind}(A_l) = \text{ind}(A'_l)$ for any finite field extension l/k . In fact, by [Ka00, Lemma 7.13] the latter is equivalent to $\langle A \rangle = \langle A' \rangle$ in $\text{Br}(k)$. In particular, if $\exp(A) = \exp(A')$ is 2, 3, 4, 6, we obtain $A \simeq A'$ or A'^{op} .

In the opposite direction, let $A \simeq A'$ or A'^{op} . By conditions (i) and (iii) one has two motivic decompositions (*) with the same sets of indices $\{i\}$ and $\{j\}$. Now according to 2.4 the motives of $\text{SB}(A)$ and $\text{SB}(A')$ are isomorphic and, hence, so are $\mathcal{M}(X)$ and $\mathcal{M}(Y)$. $\square$

The following obvious consequences of Theorem 1.4 are used in the proof of Theorem 1.1.

4.4 Corollary. *Let $X = \text{SB}_{1,\dots,n}(A)$ and $Y = \text{SB}_{1,\dots,n}(A')$ be twisted forms of the variety of complete flags of type A_n . Assume the respective central simple algebras A and A' have exponents 1, 2, 3, 4 or 6. Then*

$$\mathcal{M}(X) \simeq \mathcal{M}(Y) \Leftrightarrow X \simeq Y.$$

4.5 Corollary. *Let X and Y be twisted forms of the variety of complete flags B_2/B . Then*

$$\mathcal{M}(X) \simeq \mathcal{M}(Y) \Leftrightarrow X \simeq Y.$$

Proof. The proof repeats the proof of 1.4 observing that the motivic decompositions (*) is provided by [CPSZ, Cor. 2.9]. $\square$

4.6. Consider the pseudo-abelian completion $\mathcal{M}(G, R)$ of the category of motives of projective G -homogeneous varieties with R -coefficients, where G is a group of inner type A_n and R is a ring of coefficients. Such categories were defined and extensively studied in [CM04]. In particular, it was proven that any object of $\mathcal{M}(G, R)$, where R is a discrete valuation ring, has a unique direct sum decomposition into indecomposable objects. Modulo this result the proof of 1.4 immediately implies the following

4.7 Corollary. *Let G be an adjoint semi-simple group of inner type A_n . Let R be a ring such that any object of $\mathcal{M}(G, R)$ has a unique direct sum decomposition into indecomposable objects. Let $X = \text{SB}_{n_1, \dots, n_r}(A)$ and $Y = \text{SB}_{m_1, \dots, m_r}(A')$ be two twisted flag varieties of type A_n given by central simple algebras A and A' of prime degree. Assume that $\mathcal{M}(X_s) \simeq \mathcal{M}(Y_s)$. Then*

$$\mathcal{M}(X) \simeq \mathcal{M}(Y) \text{ in } \mathcal{M}(G, R) \iff \langle A \rangle = \langle A' \rangle \text{ in } \text{Br}(k).$$

Acknowledgements

These notes appeared as a result of the seminar organized at the University of Bielefeld in the winter of 2004-2005. We give our thanks to the participants of the seminar V.Petrov, O.Roendigs, B.Calmès and others for their remarks.

References

- [Ar82] M. Artin. Brauer-Severi varieties. In A. Dold, B. Eckmann (eds), *Lecture Notes in Mathematics*, 917, Springer-Verlag, Berlin-Heidelberg-New York, 1982, 194–210.
- [Bo03] J.-P. Bonnet. Un isomorphisme motivique entre deux variétés homogènes projectives sous l'action d'un groupe de type G_2 . *Doc. Math.* **8**, 2003, 247–277.

- [Ca72] R.W. Carter. Simple groups of Lie type. Pure and Applied Math. Eds. R. Courant, L. Bers, J.J. Stoker, vol. XXVIII, John Wiley and Sons, 1972.
- [CM04] V. Chernousov, A. Merkurjev. Motivic decomposition of projective homogeneous varieties and the Krull-Schmidt theorem. Preprint 2004.
- [CPSZ] B. Calmès, V. Petrov, N. Semenov, K. Zainoulline. Chow motives of twisted flag varieties. Preprint 2005, 28pp.
Available from <http://www.math.uni-bielefeld.de/LAG/man/186.html>
- [De77] M. Demazure. Automorphismes et déformations des variétés de Borel. *Inv. Math.*, **39** (1977), 179–186.
- [Fu98] W. Fulton. Intersection theory. Second edition, Springer-Verlag, Berlin-Heidelberg, 1998.
- [Izh98] O. Izhboldin. Motivic equivalence of quadratic forms. *Doc. Math.* **3** (1998), 341–351.
- [Inv] M.-A. Knus, A. Merkurjev, M. Rost, J.-P. Tignol. The book of involutions. AMS Colloquium Publications, vol. 44, 1998.
- [Ka96] N. Karpenko. The Grothendieck Chow-motifs of Severi-Brauer varieties. *St. Petersburg Math. J.* **7** (1996), no. 4, 649–661.
- [Ka00] N. Karpenko. Criteria of motivic equivalence for quadratic forms and central simple algebras. *Math. Ann.* **317** (2000), no. 3, 585–611.
- [Ko91] B. Köck. Chow motif and higher Chow theory of G/P . *Manuscripta Math.* **70**, 1991, 363–372.
- [Ma68] Y. Manin, Correspondences, motives and monoidal transformations. *Matematicheskij Sbornik* 77 (119) (1968), no. 4, 475–507 (in Russian). Engl. transl.: *Math. USSR Sb.* **6** (1968), 439–470.
- [Me95] A.S. Merkurjev. Zero cycles on some involution varieties. *Zap. Nauchn. Sem. POMI*, **227**, 1995, 93–105. Transl.: *J. Math. Sci.*, **89**, no. 2 (1998), 1141–1148.
- [MPW] A.S. Merkurjev, I.A. Panin, A.R. Wadsworth. Index reduction formulas for twisted flag varieties. I. *K-Theory*, **10** (1996), no. 6, 517–596.

- [NSZ] S. Nikolenko, N. Semenov, K. Zainoulline. Motivic decomposition of anisotropic varieties of type F_4 and generalized Rost motives. Preprint, 2005. Available from Arxiv Preprint Server, math.AG/0502382.
- [KR94] I. Kersten, U. Rehmann. General splitting of reductive groups. *Tohoku Math. J.* **46** (1994), 35–70.
- [Ro98] M. Rost. The motive of a Pfister form. Preprint, 1998 Available from <http://www.math.uni-bielefeld.de/~rost/>.
- [Sw89] R.G. Swan. Zero cycles on quadric hypersurfaces. *Proc. AMS*, **107**, no. 1, (1989), 43–46.
- [Ti66] J. Tits. Classification of algebraic semisimple groups. In *Algebraic Groups and Discontinuous Subgroups (Proc. Symp. Pure Math.)*, Amer. Math. Soc., Providence, R.I., 1966.
- [To04] B. Totaro. Splitting fields for E_8 -torsors. *Duke Math. J.* **121** (2004), no. 3, 425–455.
- [Vi03] A. Vishik. Motives of quadrics with applications to the theory of quadratic forms. *Lect. Notes Math.*, **1835**, 2003, 25–101.